



IMPORTANT NOTE FOR OUR SURGICAL PATIENTS

(Prior to your surgery): This *surgical reference sheet* NEEDS TO BE PLACED in an area where it can conveniently be referred to by you, after your surgery. A night-stand or refrigerator are common places many patients choose. After your surgery has occurred, it is common to have a variety of questions. This surgical reference sheet will be your guide and help answer these common questions:

Carotid Endarterectomy Surgical Patient

What to Expect After Surgery

- Some bruising.
- Some swelling.
- Normal for incision to drain small amounts of fluid for 2-3 days after surgery.
- Normal to develop hard knots around incision. May apply warm, moist heat 3-4 times per day to help with tenderness and reduce knot.

Surgical Site Care Instructions

- You may remove bandages 24 hrs after surgery.
- Steri-strips should remain on for 7 days after surgery.
- You may shower 24 hrs after surgery. Incision(s) can be washed with soap and water.

Pain Mgmt After Surgery

- You will be given an appropriate prescription for pain medicine. Take as directed by your physician.
- After that initial prescription, a patient's remaining pain is generally managed with over-the-counter pain medications. Pain medication such as ibuprofen (Advil™ or Motrin™) **OR** acetaminophen (Tylenol™) **OR** naproxen (Aleve™) can be taken. Follow label instructions.

Other Helpful Information

- May drive once you are no longer using prescription pain medicine.
- Avoid strenuous activity for 2 weeks after surgery. Then proceed with activities as tolerated.
- Avoid alcoholic beverages and tobacco use.

When to Contact Our Office

- Fever of 101° or greater.
- Increased redness or swelling around incision.
- Have a sudden or large amount of drainage from incision.
- Have a sudden increase in pain.
- Experience difficulty swallowing.